

Recent Results of Prenatal Diagnosis in the Czech Republic

**Antonin Sipek jr.¹, Vladimir Gregor^{2,3}, Antonin Sipek^{2,3}
Jiri Horacek⁴ and Pavel Langhammer⁵**

1 – Institute of Biology and Medical Genetics, Charles University, 1st Faculty of Medicine and General University Hospital, Prague, Czech Republic

2 – Department of Medical Genetics, Thomayer Hospital, Prague, Czech Republic

3 – Screening centre, Pronatal Sanatory, Prague, Czech Republic

4 – Gennet Ltd, Prague, Czech Republic

5 – Institute for Health Information and Statistics of the Czech Republic



<http://www.vrozene-vady.cz/>

History and Current State I

National Registry of Congenital Anomalies of the Czech Republic (NRCA)

- Unofficial monitoring in former Czechoslovakia started in **1961**
- Official monitoring started on 1st of January **1964**
- **First stage (1964 – 1974)** – only **36** selected diagnoses of congenital anomalies (CA) were registered
- **Second stage (1975 – 1993)** – **60** diagnoses of CA registered
- **Present time: (1994 – now)** – **all cases** in terminations of pregnancies (TOPs), stillbirths and live births are registered (age limit for reporting = **15 years**)

Cooperating centre:



eurocat
european surveillance of
congenital anomalies

History and Current State II

Registry: population based (whole area of the Czech Republic)

Law: The registration is compulsory, required by the Internal Law of Ministry of Health (nr. 14/2001). The database is run by the Institute of Health Information and Statistics of the Czech Republic


Cases: all cases in TOPs, live births and stillbirths (above 1000g) are reported

Coding: ICD-10 (international), no verbal description


Sources: Multiple sources, including departments of medical genetics, genetic laboratories, pediatric and neonatology departments, delivery units, ultrasound diagnostics departments etc.

Termination of Pregnancy: Legal, up to the 24th week of gestation (from genetic reasons)


Prenatal diagnosis - Numbers



Invasive prenatal diagnosis




Effectiveness




Number of invasive procedures needed for one diagnosis of Down syndrome

Prenatal diagnosis - Indications




Spectrum of diagnoses



- CA of neural tissue
- CA of cardiovascular tract
- Cleft lip and/or palate
- CA of urinary tract
- Diaphragmatic hernia and abdominal wall defects
- CA of bone and muscle tissue
- Cromosomal abnormalities
- Other

Spectrum of diagnoses – Prenatal diagnosis – All cases

Maternal age – Time trends




Anencephaly

per 10 000 live births


■ Births

■ Prenatal diagnosis




Anencephaly – All cases in live births and prenatal diagnosis

Anencephaly




Anencephaly – relative number of prenatally diagnosed cases (%)

Spina bifida



Spina bifida – All cases in live births and prenatal diagnosis

Spina bifida




Spina bifida – relative number of prenatally diagnosed cases (%)

Omphalocele

per 10 000 live births


■ Births

■ Prenatal diagnosis




Omphalocele – All cases in live births and prenatal diagnosis

Omphalocele




Omphalocele – relative number of prenatally diagnosed cases (%)

Gastroschisis



Gastroschisis – All cases in live births and prenatal diagnosis

Gastroschisis




Gastroschisis – relative number of prenatally diagnosed cases (%)

Down syndrome

Per 10 000 live births


■ Births

■ Prenatal diagnosis




Down syndrome – All cases in live births and prenatal diagnosis

Down syndrome



Down syndrome – relative number of prenatally diagnosed cases (%)

Down syndrome - ICBDSR



<http://www.icbdsr.org>

Down syndrome – relative number of prenatally diagnosed cases (%)

Conclusions

1. *The number of prenatally diagnosed cases of congenital anomalies is increasing*
2. *The number of congenital anomalies in live births is decreasing*
3. *The indication criteria for the invasive prenatal diagnosis are changing during last years*
4. *The average maternal age is notably increasing during last years*
5. *The early prenatal diagnosis (during the first trimester of gravidity) is becoming much more important during last years*

Acknowledgements

This analysis would not be possible without the cooperation of all helpful workers (geneticists, pediatricians, gynecologists and many others) who are involved in the registration process and provide the data, necessary for the surveillance of the congenital anomalies in the Czech Republic

Thank you for your attention!



antonin.sipek@lf1.cuni.cz

<http://www.vrozene-vady.cz/>